



1325 4th Avenue, Suite 1230 Seattle,
Washington 98101
TEL: 206.624.1773
FAX: 800.886.1511
info@seattlesmilesdental.com
seattlesmilesdental.com

Jenny Nguyen DDS | Asmeret Tesfahun DDS

INFANT FRENECTOMY (UNDER AGE 1) EVALUATION REQUEST FORM

Please return form to us by email (preferred) or fax

Today's date: _____ Examination date: _____

Referred from (provider): _____ Specialty/field: _____

Referral address: _____

Referral phone number: _____ Referral email: _____

Patient's name: _____ DOB: _____

Parent(s) or guardian: _____

Patient address: _____

Contact phone number and name: _____

Should we contact the patient? ☐ Yes ☐ No

<p>Reason for referral (check all that apply):</p> <ul style="list-style-type: none"><input type="checkbox"/> Evaluation and report only<input type="checkbox"/> Evaluation and treatment as needed<input type="checkbox"/> Lingual (tongue) frenectomy<input type="checkbox"/> Labial (lip) frenectomy<ul style="list-style-type: none"><input type="checkbox"/> Maxillary<input type="checkbox"/> Mandibular<input type="checkbox"/> Buccal (cheek) frenectomy Location(s) (UR/LR/UL/LL): _____ <p>Family history of lip/tongue-tie? <input type="checkbox"/> Yes <input type="checkbox"/> No</p> <p>Has the baby had any of the following? (If so, referral to OT/PT may be recommended after the evaluation but prior to treatment):</p> <ul style="list-style-type: none"><input type="checkbox"/> signs of aspiration (choking, coughing, respiratory infections, fevers, tearing of eyes, color change)<input type="checkbox"/> underlying neuromuscular issue or medical diagnosis (e.g., down syndrome, cleft, seizure, etc.)<input type="checkbox"/> oral aversion (extensive gagging, combative at sight, touch or smells near mouth)<input type="checkbox"/> torticollis (twisted neck)<input type="checkbox"/> plagiocephaly (flat head)	<p>Baby's signs/symptoms:</p> <ul style="list-style-type: none"><input type="checkbox"/> poor latch<input type="checkbox"/> falls asleep while attempting to nurse<input type="checkbox"/> slides off the nipple when attempting to latch<input type="checkbox"/> colic symptoms<input type="checkbox"/> reflux symptoms (spitting up, vomiting, stuffy nose)<input type="checkbox"/> poor weight gain<input type="checkbox"/> gumming or chewing of nipple when nursing<input type="checkbox"/> unable to hold a pacifier in his or her mouth<input type="checkbox"/> mouth breathing/open mouth resting posture<input type="checkbox"/> snoring/noisy breathing<input type="checkbox"/> short sleep episodes requiring feeding every 2-3 hours <p>Mother's signs/symptoms:</p> <ul style="list-style-type: none"><input type="checkbox"/> creased/flattened/blanched nipples after nursing<input type="checkbox"/> cracked/bruised/blistered nipples<input type="checkbox"/> bleeding nipples<input type="checkbox"/> severe pain when infant attempts to latch<input type="checkbox"/> poor/incomplete breast drainage<input type="checkbox"/> infected nipples or breasts<input type="checkbox"/> plugged ducts<input type="checkbox"/> mastitis or nipple thrush<input type="checkbox"/> use of nipple shield
--	--

Additional notes or concerns: